Exhibit 24

United States Environmental Protection Agency

April 2011

Inventory of U.S. Greenhouse Gas Emissions and Sinks: 1990–2009

Executive Summary

An emissions inventory that identifies and quantifies a country's primary anthropogenic[1] sources and sinks of greenhouse gases is essential for addressing climate change. This inventory adheres to both (1) a comprehensive and detailed set of methodologies for estimating sources and sinks of anthropogenic greenhouse gases, and (2) a common and consistent mechanism that enables Parties to the United Nations Framework Convention on Climate Change (UNFCCC) to compare the relative contribution of different emission sources and greenhouse gases to climate change.

In 1992, the United States signed and ratified the UNFCCC. As stated in Article 2 of the UNFCCC, "The ultimate objective of this Convention and any related legal instruments that the Conference of the Parties may adopt is to achieve, in accordance with the relevant provisions of the Convention, stabilization of greenhouse gas concentrations in the atmosphere at a level that would prevent dangerous anthropogenic interference with the climate system. Such a level should be achieved within a time-frame sufficient to allow ecosystems to adapt naturally to climate change, to ensure that food production is not threatened and to enable economic development to proceed in a sustainable manner."[2]

All material taken from the *Inventory of U.S. Greenhouse Gas Emissions and Sinks: 1990–2009*, U.S. Environmental Protection Agency, Office of Atmospheric Programs, EPA 430-R-11-005, April 2011. You may electronically download the full inventory report from U.S. EPA's Global Climate Change web page at: **www.epa.gov/climatechange/emissions/usinventory.html.**

Parties to the Convention, by ratifying, "shall develop, periodically update, publish and make available…national inventories of anthropogenic emissions by sources and removals by sinks of all greenhouse gases not controlled by the *Montreal Protocol*, using comparable methodologies…"[3] The United States views the Inventory as an opportunity to fulfill these commitments.

This chapter summarizes the latest information on U.S. anthropogenic greenhouse gas emission trends from 1990 through 2009. To ensure that the U.S. emission inventory is comparable to those of other UNFCCC Parties, the estimates presented here were calculated using methodologies consistent with those recommended in the Revised 1996 Intergovernmental Panel on Climate Change (IPCC) *Guidelines for National Greenhouse Gas Inventories* (IPCC/UNEP/OECD/IEA 1997), the IPCC *Good Practice Guidance and Uncertainty Management in National Greenhouse Gas Inventories* (IPCC 2000), and the IPCC *Good Practice Guidance for Land Use, Land-Use Change, and Forestry* (IPCC 2003). Additionally, the U.S. emission inventory has continued to incorporate new methodologies and data from the *2006 IPCC Guidelines for National Greenhouse Gas Inventories* (IPCC 2006). The structure of the inventory report is consistent

[1] The term "anthropogenic", in this context, refers to greenhouse gas emissions and removals that are a direct result of human activities or are the result of natural processes that have been affected by human activities (IPCC/UNEP/OECD/IEA 1997).

[2] Article 2 of the Framework Convention on Climate Change published by the UNEP/WMO Information Unit on Climate Change. See <http://unfccc.int>.

[3] Article 4(1)(a) of the United Nations Framework Convention on Climate Change (also identified in Article 12). Subsequent decisions by the Conference of the Parties elaborated the role of Annex I Parties in preparing national inventories. See <http://unfccc.int>.

with the UNFCCC guidelines for inventory reporting.[4] For most source categories, the IPCC methodologies were expanded, resulting in a more comprehensive and detailed estimate of emissions.

Box ES-1: Methodological approach for estimating and reporting U.S. emissions and sinks

In following the UNFCCC requirement under Article 4.1 to develop and submit national greenhouse gas emissions inventories, the emissions and sinks presented in the inventory report are organized by source and sink categories and calculated using internationally-accepted methods provided by the IPCC.[5] Additionally, the calculated emissions and sinks in a given year for the U.S. are presented in a common manner in line with the UNFCCC reporting guidelines for the reporting of inventories under this international agreement.[6] The use of consistent methods to calculate emissions and sinks by all nations providing their inventories to the UNFCCC ensures that these reports are comparable. In this regard, U.S. emissions and sinks reported in this inventory report are comparable to emissions and sinks reported by other countries. Emissions and sinks provided in this inventory do not preclude alternative examinations, but rather this inventory report presents emissions and sinks in a common format consistent with how countries are to report inventories under the UNFCCC. The inventory report itself follows this standardized format, and provides an explanation of the IPCC methods used to calculate emissions and sinks, and the manner in which those calculations are conducted.

ES.1. Background Information

Naturally occurring greenhouse gases include water vapor, carbon dioxide (CO_2), methane (CH_4), nitrous oxide (N_2O), and ozone (O_3). Several classes of halogenated substances that contain fluorine, chlorine, or bromine are also greenhouse gases, but they are, for the most part, solely a product of industrial activities. Chlorofluorocarbons (CFCs) and hydrochlorofluorocarbons (HCFCs) are halocarbons that contain chlorine, while halocarbons that contain bromine are referred to as bromofluorocarbons (i.e., halons). As stratospheric ozone depleting substances, CFCs, HCFCs, and halons are covered under the *Montreal Protocol* on Substances that Deplete the Ozone Layer. The UNFCCC defers to this earlier international treaty. Consequently, Parties to the UNFCCC are not required to include these gases in their national greenhouse gas emission inventories.[7] Some other fluorine-containing halogenated substances—hydrofluorocarbons (HFCs), perfluorocarbons (PFCs), and sulfur hexafluoride (SF_6)—do not deplete stratospheric ozone but are potent greenhouse gases. These latter substances are addressed by the UNFCCC and accounted for in national greenhouse gas emission inventories.

There are also several gases that do not have a direct global warming effect but indirectly affect terrestrial and/or solar radiation absorption by influencing the formation or destruction of greenhouse gases, including tropospheric and stratospheric ozone. These gases include carbon monoxide (CO), oxides of nitrogen (NO_x), and non-CH_4 volatile organic compounds (NMVOCs). Aerosols, which are extremely small particles or liquid droplets, such as those produced by sulfur dioxide (SO_2) or elemental carbon emissions, can also affect the absorptive characteristics of the atmosphere.

Although the direct greenhouse gases CO_2, CH_4, and N_2O occur naturally in the atmosphere, human activities have changed their atmospheric concentrations. From the pre-industrial era (i.e., ending about 1750) to 2005, concentrations of these greenhouse gases have increased globally by 36, 148, and 18 percent, respectively (IPCC 2007).

[4] See < http://unfccc.int/resource/docs/2006/sbsta/eng/09.pdf>.

[5] See < http://www.ipcc-nggip.iges.or.jp/public/index.html>.

[6] See < http://unfccc.int/national_reports/annex_i_ghg_inventories/national_inventories_submissions/items/5270.php>.

[7] Emissions estimates of CFCs, HCFCs, halons and other ozone-depleting substances are included in the annexes of the inventory report for informational purposes.

Beginning in the 1950s, the use of CFCs and other stratospheric ozone depleting substances (ODS) increased by nearly 10 percent per year until the mid-1980s, when international concern about ozone depletion led to the entry into force of the *Montreal Protocol*. Since then, the production of ODS is being phased out. In recent years, use of ODS substitutes such as HFCs and PFCs has grown as they begin to be phased in as replacements for CFCs and HCFCs. Accordingly, atmospheric concentrations of these substitutes have been growing (IPCC 2007).

Global Warming Potentials

Gases in the atmosphere can contribute to the greenhouse effect both directly and indirectly. Direct effects occur when the gas itself absorbs radiation. Indirect radiative forcing occurs when chemical transformations of the substance produce other greenhouse gases, when a gas influences the atmospheric lifetimes of other gases, and/or when a gas affects atmospheric processes that alter the radiative balance of the earth (e.g., affect cloud formation or albedo).[8] The IPCC developed the global warming potential (GWP) concept to compare the ability of each greenhouse gas to trap heat in the atmosphere relative to another gas.

The GWP of a greenhouse gas is defined as the ratio of the time-integrated radiative forcing from the instantaneous release of 1 kilogram (kg) of a trace substance relative to that of 1 kg of a reference gas (IPCC 2001). Direct radiative effects occur when the gas itself is a greenhouse gas. The reference gas used is CO_2, and therefore GWP-weighted emissions are measured in teragrams (or million metric tons) of CO_2 equivalent (Tg CO_2 Eq.).[9, 10] All gases in this Executive Summary are presented in units of Tg CO_2 Eq.

The UNFCCC reporting guidelines for national inventories were updated in 2006,[11] but continue to require the use of GWPs from the IPCC Second Assessment Report (SAR) (IPCC 1996). This requirement ensures that current estimates of aggregate greenhouse gas emissions for 1990 to 2009 are consistent with estimates developed prior to the publication of the IPCC Third Assessment Report (TAR) (IPCC 2001) and the IPCC Fourth Assessment Report (AR4) (IPCC 2007). Therefore, to comply with international reporting standards under the UNFCCC, official emission estimates are reported by the United States using SAR GWP values. All estimates are provided throughout the inventory report in both CO_2 equivalents and unweighted units. A comparison of emission values using the SAR GWPs versus the TAR and AR4 GWPs can be found in Chapter 1 and, in more detail, in Annex 6.1 of the inventory report. The GWP values used in the inventory report are listed below in Table ES-1.

Table ES-1: Global Warming Potentials (100-Year Time Horizon) Used in the Inventory Report

Gas	GWP
CO_2	1
CH_4*	21
N_2O	310
HFC-23	11,700
HFC-32	650
HFC-125	2,800
HFC-134a	1,300
HFC-143a	3,800
HFC-152a	140
HFC-227ea	2,900
HFC-236fa	6,300
HFC-4310mee	1,300
CF_4	6,500
C_2F_6	9,200
C_4F_{10}	7,000
C_6F_{14}	7,400
SF_6	23,900

Source: IPCC (1996)

* The CH_4 GWP includes the direct effects and those indirect effects due to the production of tropospheric ozone and stratospheric water vapor. The indirect effect due to the production of CO_2 is not included.

[8] Albedo is a measure of the earth's reflectivity, and is defined as the fraction of the total solar radiation incident on a body that is reflected by it.

[9] Carbon comprises 12/44^{ths} of carbon dioxide by weight.

[10] One teragram is equal to 10^{12} grams or one million metric tons.

[11] See <http://unfccc.int/resource/docs/2006/sbsta/eng/09.pdf>.

Global warming potentials are not provided for CO, NO_x, NMVOCs, SO_2, and aerosols because there is no agreed-upon method to estimate the contribution of gases that are short-lived in the atmosphere, spatially variable, or have only indirect effects on radiative forcing (IPCC 1996).

ES.2. Recent Trends in U.S. Greenhouse Gas Emissions and Sinks

In 2009, total U.S. greenhouse gas emissions were 6,633.2 Tg or million metric tons CO_2 Eq. While total U.S. emissions have increased by 7.3 percent from 1990 to 2009, emissions decreased from 2008 to 2009 by 6.1 percent (427.9 Tg CO_2 Eq.). This decrease was primarily due to (1) a decrease in economic output resulting in a decrease in energy consumption across all sectors; and (2) a decrease in the carbon intensity of fuels used to generate electricity due to fuel switching as the price of coal increased, and the price of natural gas decreased significantly. Since 1990, U.S. emissions have increased at an average annual rate of 0.4 percent.

Figure ES-1 through Figure ES-3 illustrate the overall trends in total U.S. emissions by gas, annual changes, and absolute change since 1990.

Table ES-2 provides a detailed summary of U.S. greenhouse gas emissions and sinks for 1990 through 2009.

Figure ES-1



Figure ES-2



Figure ES-3



Table ES-2: Recent Trends in U.S. Greenhouse Gas Emissions and Sinks (Tg or million metric tons CO_2 Eq.)

Gas/Source	1990	2000	2005	2006	2007	2008	2009
CO_2	**5,099.7**	**5,975.0**	**6,113.8**	**6,021.1**	**6,120.0**	**5,921.4**	**5,505.2**
Fossil Fuel Combustion	4,738.4	5,594.8	5,753.2	5,653.1	5,756.7	5,565.9	5,209.0
Electricity Generation	1,820.8	2,296.9	2,402.1	2,346.4	2,412.8	2,360.9	2,154.0
Transportation	1,485.9	1,809.5	1,896.6	1,878.1	1,894.0	1,789.9	1,719.7
Industrial	846.5	851.1	823.1	848.2	842.0	802.9	730.4
Residential	338.3	370.7	357.9	321.5	342.4	348.2	339.2
Commercial	219.0	230.8	223.5	208.6	219.4	224.2	224.0
U.S. Territories	27.9	35.9	50.0	50.3	46.1	39.8	41.7
Non-Energy Use of Fuels	118.6	144.9	143.4	145.6	137.2	141.0	123.4
Iron and Steel Production & Metallurgical Coke Production	99.5	85.9	65.9	68.8	71.0	66.0	41.9
Natural Gas Systems	37.6	29.9	29.9	30.8	31.1	32.8	32.2
Cement Production	33.3	40.4	45.2	45.8	44.5	40.5	29.0
Incineration of Waste	8.0	11.1	12.5	12.5	12.7	12.2	12.3
Ammonia Production and Urea Consumption	16.8	16.4	12.8	12.3	14.0	11.9	11.8
Lime Production	11.5	14.1	14.4	15.1	14.6	14.3	11.2
Cropland Remaining Cropland	7.1	7.5	7.9	7.9	8.2	8.7	7.8
Limestone and Dolomite Use	5.1	5.1	6.8	8.0	7.7	6.3	7.6
Soda Ash Production and Consumption	4.1	4.2	4.2	4.2	4.1	4.1	4.3
Aluminum Production	6.8	6.1	4.1	3.8	4.3	4.5	3.0
Petrochemical Production	3.3	4.5	4.2	3.8	3.9	3.4	2.7
Carbon Dioxide Consumption	1.4	1.4	1.3	1.7	1.9	1.8	1.8
Titanium Dioxide Production	1.2	1.8	1.8	1.8	1.9	1.8	1.5
Ferroalloy Production	2.2	1.9	1.4	1.5	1.6	1.6	1.5
Wetlands Remaining Wetlands	1.0	1.2	1.1	0.9	1.0	1.0	1.1
Phosphoric Acid Production	1.5	1.4	1.4	1.2	1.2	1.2	1.0
Zinc Production	0.7	1.0	1.1	1.1	1.1	1.2	1.0
Lead Production	0.5	0.6	0.6	0.6	0.6	0.6	0.5
Petroleum Systems	0.6	0.5	0.5	0.5	0.5	0.5	0.5
Silicon Carbide Production and Consumption	0.4	0.2	0.2	0.2	0.2	0.2	0.1
Land Use, Land-Use Change, and Forestry (Sink) [a]	*(861.5)*	*(576.6)*	*(1,056.5)*	*(1,064.3)*	*(1,060.9)*	*(1,040.5)*	*(1,015.1)*
Biomass – Wood [b]	*215.2*	*218.1*	*206.9*	*203.8*	*203.3*	*198.4*	*183.8*
International Bunker Fuels [c]	*111.8*	*98.5*	*109.7*	*128.4*	*127.6*	*133.7*	*123.1*
Biomass – Ethanol [b]	*4.2*	*9.4*	*23.0*	*31.0*	*38.9*	*54.8*	*61.2*
CH_4	**674.9**	**659.9**	**631.4**	**672.1**	**664.6**	**676.7**	**686.3**
Natural Gas Systems	189.8	209.3	190.4	217.7	205.2	211.8	221.2
Enteric Fermentation	132.1	136.5	136.5	138.8	141.0	140.6	139.8
Landfills	147.4	111.7	112.5	111.7	111.3	115.9	117.5
Coal Mining	84.1	60.4	56.9	58.2	57.9	67.1	71.0
Manure Management	31.7	42.4	46.6	46.7	50.7	49.4	49.5
Petroleum Systems	35.4	31.5	29.4	29.4	30.0	30.2	30.9
Wastewater Treatment	23.5	25.2	24.3	24.5	24.4	24.5	24.5
Forest Land Remaining Forest Land	3.2	14.3	9.8	21.6	20.0	11.9	7.8
Rice Cultivation	7.1	7.5	6.8	5.9	6.2	7.2	7.3
Stationary Combustion	7.4	6.6	6.6	6.2	6.5	6.5	6.2
Abandoned Underground Coal Mines	6.0	7.4	5.5	5.5	5.6	5.9	5.5
Mobile Combustion	4.7	3.4	2.5	2.3	2.2	2.0	2.0
Composting	0.3	1.3	1.6	1.6	1.7	1.7	1.7
Petrochemical Production	0.9	1.2	1.1	1.0	1.0	0.9	0.8
Iron and Steel Production & Metallurgical Coke Production	1.0	0.9	0.7	0.7	0.7	0.6	0.4
Field Burning of Agricultural Residues	0.3	0.3	0.2	0.2	0.2	0.3	0.2
Ferroalloy Production	+	+	+	+	+	+	+

Table ES-2: Recent Trends in U.S. Greenhouse Gas Emissions and Sinks (Tg or million metric tons CO_2 Eq.) (continued)

Gas/Source	1990	2000	2005	2006	2007	2008	2009
Silicon Carbide Production and Consumption	+	+	+	+	+	+	+
Incineration of Waste	+	+	+	+	+	+	+
International Bunker Fuels[c]	*0.2*	*0.1*	*0.1*	*0.2*	*0.2*	*0.2*	*0.1*
N_2O	**315.2**	**341.0**	**322.9**	**326.4**	**325.1**	**310.8**	**295.6**
Agricultural Soil Management	197.8	206.8	211.3	208.9	209.4	210.7	204.6
Mobile Combustion	43.9	53.2	36.9	33.6	30.3	26.1	23.9
Manure Management	14.5	17.1	17.3	18.0	18.1	17.9	17.9
Nitric Acid Production	17.7	19.4	16.5	16.2	19.2	16.4	14.6
Stationary Combustion	12.8	14.6	14.7	14.4	14.6	14.2	12.8
Forest Land Remaining Forest Land	2.7	12.1	8.4	18.0	16.7	10.1	6.7
Wastewater Treatment	3.7	4.5	4.8	4.8	4.9	5.0	5.0
N_2O from Product Uses	4.4	4.9	4.4	4.4	4.4	4.4	4.4
Adipic Acid Production	15.8	5.5	5.0	4.3	3.7	2.0	1.9
Composting	0.4	1.4	1.7	1.8	1.8	1.9	1.8
Settlements Remaining Settlements	1.0	1.1	1.5	1.5	1.6	1.5	1.5
Incineration of Waste	0.5	0.4	0.4	0.4	0.4	0.4	0.4
Field Burning of Agricultural Residues	0.1	0.1	0.1	0.1	0.1	0.1	0.1
Wetlands Remaining Wetlands	+	+	+	+	+	+	+
International Bunker Fuels [c]	*1.1*	*0.9*	*1.0*	*1.2*	*1.2*	*1.2*	*1.1*
HFCs	**36.9**	**103.2**	**120.2**	**123.5**	**129.5**	**129.4**	**125.7**
Substitution of Ozone Depleting Substances [d]	0.3	74.3	104.2	109.4	112.3	115.5	120.0
HCFC-22 Production	36.4	28.6	15.8	13.8	17.0	13.6	5.4
Semiconductor Manufacture	0.2	0.3	0.2	0.3	0.3	0.3	0.3
PFCs	**20.8**	**13.5**	**6.2**	**6.0**	**7.5**	**6.6**	**5.6**
Semiconductor Manufacture	2.2	4.9	3.2	3.5	3.7	4.0	4.0
Aluminum Production	18.5	8.6	3.0	2.5	3.8	2.7	1.6
SF_6	**34.4**	**20.1**	**19.0**	**17.9**	**16.7**	**16.1**	**14.8**
Electrical Transmission and Distribution	28.4	16.0	15.1	14.1	13.2	13.3	12.8
Magnesium Production and Processing	5.4	3.0	2.9	2.9	2.6	1.9	1.1
Semiconductor Manufacture	0.5	1.1	1.0	1.0	0.8	0.9	1.0
Total	**6,181.8**	**7,112.7**	**7,213.5**	**7,166.9**	**7,263.4**	**7,061.1**	**6,633.2**
Net Emissions (Sources and Sinks)	**5,320.3**	**6,536.1**	**6,157.1**	**6,102.6**	**6,202.5**	**6,020.7**	**5,618.2**

+ Does not exceed 0.05 Tg CO_2 Eq.
[a] Parentheses indicate negative values or sequestration. The net CO_2 flux total includes both emissions and sequestration, and constitutes a net sink in the United States. Sinks are only included in net emissions total.
[b] Emissions from Wood Biomass and Ethanol Consumption are not included specifically in summing energy sector totals. Net carbon fluxes from changes in biogenic carbon reservoirs are accounted for in the estimates for Land Use, Land-Use Change, and Forestry.
[c] Emissions from International Bunker Fuels are not included in totals.
[d] Small amounts of PFC emissions also result from this source.
Note: Totals may not sum due to independent rounding.

Figure ES-4 illustrates the relative contribution of the direct greenhouse gases to total U.S. emissions in 2009. The primary greenhouse gas emitted by human activities in the United States was CO_2, representing approximately 83.0 percent of total greenhouse gas emissions. The largest source of CO_2, and of overall greenhouse gas emissions, was fossil fuel combustion. Methane emissions, which have increased by 1.7 percent since 1990, resulted primarily from natural gas systems, enteric fermentation associated with domestic livestock, and decomposition of wastes in landfills. Agricultural soil management and mobile source fuel combustion were the major sources of N_2O emissions. Ozone depleting substance substitute emissions and emissions of HFC-23 during the production of HCFC-22 were the primary contributors to aggregate HFC emissions. PFC emissions resulted as a byproduct of primary aluminum production and from semiconductor manufacturing, while electrical transmission and distribution systems accounted for most SF_6 emissions.

Overall, from 1990 to 2009, total emissions of CO_2 and CH_4 increased by 405.5 Tg CO_2 Eq. (8.0 percent) and 11.4 Tg CO_2 Eq. (1.7 percent), respectively. Conversely, N_2O emissions decreased by 19.6 Tg CO_2 Eq. (6.2 percent). During the same period, aggregate weighted emissions of HFCs, PFCs, and SF_6 rose by 54.1 Tg CO_2 Eq. (58.8 percent). From 1990 to 2009, HFCs increased by 88.8 Tg CO_2 Eq. (240.41 percent), PFCs decreased by 15.1 Tg CO_2 Eq. (73.0 percent), and SF_6 decreased by 19.5 Tg CO_2 Eq. (56.8 percent). Despite being emitted in smaller quantities relative to the other principal greenhouse gases, emissions of HFCs, PFCs, and SF_6 are significant because many of these gases have extremely high global warming potentials and, in the cases of PFCs and SF_6, long atmospheric lifetimes. Conversely, U.S. greenhouse gas emissions were partly offset by carbon sequestration in forests, trees in urban areas, agricultural soils, and landfilled yard trimmings and food scraps, which, in aggregate, offset 15.3 percent of total emissions in 2009. The following sections describe each gas' contribution to total U.S. greenhouse gas emissions in more detail.

Figure ES-4

2009 Greenhouse Gas Emissions by Gas (percents based on Tg CO_2 Eq.)



Carbon Dioxide Emissions

The global carbon cycle is made up of large carbon flows and reservoirs. Billions of tons of carbon in the form of CO_2 are absorbed by oceans and living biomass (i.e., sinks) and are emitted to the atmosphere annually through natural processes (i.e., sources). When in equilibrium, carbon fluxes among these various reservoirs are roughly balanced. Since the Industrial Revolution (i.e., about 1750), global atmospheric concentrations of CO_2 have risen about 36 percent (IPCC 2007), principally due to the combustion of fossil fuels. Within the United States, fossil fuel combustion accounted for 94.6 percent of CO_2 emissions in 2009. Globally, approximately 30,313 Tg of CO_2 were added to the atmosphere through the combustion of fossil fuels in 2009, of which the United States accounted for about 18 percent.[12] Changes in land use and forestry practices can also emit CO_2 (e.g., through conversion of forest land to agricultural or urban use) or can act as a sink for CO_2 (e.g., through net additions to forest biomass). In addition to fossil-fuel combustion, several other sources emit significant quantities of CO_2. These sources include, but are not limited to non-energy use of fuels, iron and steel production and cement production (Figure ES-5).

As the largest source of U.S. greenhouse gas emissions, CO_2 from fossil fuel combustion has accounted for approximately 78 percent of GWP-weighted emissions since 1990, growing slowly from 77 percent of total GWP-weighted emissions in 1990 to 79 percent in 2009. Emissions of CO_2 from fossil fuel combustion increased at an average annual rate of 0.4 percent from 1990 to 2009. The fundamental factors influencing this trend include: (1) a generally growing domestic economy over the last 20 years, and (2) overall growth in emissions from electricity generation and transportation activities. Between 1990 and 2009, CO_2 emissions from fossil fuel combustion increased from 4,738.4 Tg CO_2 Eq. to 5,209.0 Tg CO_2 Eq.—a 9.9 percent total increase over the twenty-year period. From 2008 to 2009, these emissions decreased by 356.9 Tg CO_2 Eq. (6.4 percent), the largest decrease in any year over the twenty-year period.

[12] Global CO_2 emissions from fossil fuel combustion were taken from Energy Information Administration *International Energy Statistics 2010* <http://tonto.eia.doe.gov/cfapps/ipdbproject/IEDIndex3.cfm> EIA (2010a).

Historically, changes in emissions from fossil fuel combustion have been the dominant factor affecting U.S. emission trends. Changes in CO_2 emissions from fossil fuel combustion are influenced by many long-term and short-term factors, including population and economic growth, energy price fluctuations, technological changes, and seasonal temperatures. In the short term, the overall consumption of fossil fuels in the United States fluctuates primarily in response to changes in general economic conditions, energy prices, weather, and the availability of non-fossil alternatives. For example, in a year with increased consumption of goods and services, low fuel prices, severe summer and winter weather conditions, nuclear plant closures, and lower precipitation feeding hydroelectric dams, there would likely be proportionally greater fossil fuel consumption than a year with poor economic performance, high fuel prices, mild temperatures, and increased output from nuclear and hydroelectric plants. In the long term, energy consumption patterns respond to changes that affect the scale of consumption (e.g., population, number of cars, and size of houses), the efficiency with which energy is used in equipment (e.g., cars, power plants, steel mills, and light bulbs) and behavioral choices (e.g., walking, bicycling, or telecommuting to work instead of driving).

Figure ES-5



The five major fuel consuming sectors contributing to CO_2 emissions from fossil fuel combustion are electricity generation, transportation, industrial, residential, and commercial. Carbon dioxide emissions are produced by the electricity generation sector as they consume fossil fuel to provide electricity to one of the other four sectors, or "end-use" sectors. For the discussion below, electricity generation emissions have been distributed to each end-use sector on the basis of each sector's share of aggregate electricity consumption. This method of distributing emissions assumes that each end-use sector consumes electricity that is generated from the national average mix of fuels according to their carbon intensity. Emissions from electricity generation are also addressed separately after the end-use sectors have been discussed.

Note that emissions from U.S. territories are calculated separately due to a lack of specific consumption data for the individual end-use sectors.

Figure ES-6, Figure ES-7, and Table ES-3 summarize CO_2 emissions from fossil fuel combustion by end-use sector.

Figure ES-6

2009 CO_2 Emissions from Fossil Fuel Combustion by Sector and Fuel Type



Note: Electricity generation also includes emissions of less than 0.5 Tg CO_2 Eq. from geothermal-based electricity generation.

Figure ES-7

2009 End-Use Sector Emissions of CO_2, CH_4, and N_2O from Fossil Fuel Combustion



Table ES-3: CO_2 Emissions from Fossil Fuel Combustion by Fuel Consuming End-Use Sector (Tg or million metric tons CO_2 Eq.)

End-Use Sector	1990	2000	2005	2006	2007	2008	2009
Transportation	**1,489.0**	**1,813.0**	**1,901.3**	**1,882.6**	**1,899.0**	**1,794.6**	**1,724.1**
Combustion	1,485.9	1,809.5	1,896.6	1,878.1	1,894.0	1,789.9	1,719.7
Electricity	3.0	3.4	4.7	4.5	5.0	4.7	4.4
Industrial	**1,533.2**	**1,640.8**	**1,560.0**	**1,560.2**	**1,572.0**	**1,517.7**	**1,333.7**
Combustion	846.5	851.1	823.1	848.2	842.0	802.9	730.4
Electricity	686.7	789.8	737.0	712.0	730.0	714.8	603.3
Residential	**931.4**	**1,133.1**	**1,214.7**	**1,152.4**	**1,198.5**	**1,182.2**	**1,123.8**
Combustion	338.3	370.7	357.9	321.5	342.4	348.2	339.2
Electricity	593.0	762.4	856.7	830.8	856.1	834.0	784.6
Commercial	**757.0**	**972.1**	**1,027.2**	**1,007.6**	**1,041.1**	**1,031.6**	**985.7**
Combustion	219.0	230.8	223.5	208.6	219.4	224.2	224.0
Electricity	538.0	741.3	803.7	799.0	821.7	807.4	761.7
U.S. Territories [a]	**27.9**	**35.9**	**50.0**	**50.3**	**46.1**	**39.8**	**41.7**
Total	**4,738.4**	**5,594.8**	**5,753.2**	**5,653.1**	**5,756.7**	**5,565.9**	**5,209.0**
Electricity Generation	**1,820.8**	**2,296.9**	**2,402.1**	**2,346.4**	**2,412.8**	**2,360.9**	**2,154.0**

[a] Fuel consumption by U.S. territories (i.e., American Samoa, Guam, Puerto Rico, U.S. Virgin Islands, Wake Island, and other U.S. Pacific Islands) is included in the inventory report.
Note: Totals may not sum due to independent rounding. Combustion-related emissions from electricity generation are allocated based on aggregate national electricity consumption by each end-use sector.

Transportation End-Use Sector. Transportation activities (excluding international bunker fuels) accounted for 33 percent of CO_2 emissions from fossil fuel combustion in 2009.[13] Virtually all of the energy consumed in this end-use sector came from petroleum products. Nearly 65 percent of the emissions resulted from gasoline consumption for personal vehicle use. The remaining emissions came from other transportation activities, including the combustion of diesel fuel in heavy-

[13] If emissions from international bunker fuels are included, the transportation end-use sector accounted for 35 percent of U.S. emissions from fossil fuel combustion in 2009.

duty vehicles and jet fuel in aircraft. From 1990 to 2009, transportation emissions rose by 16 percent due, in large part, to increased demand for travel and the stagnation of fuel efficiency across the U.S. vehicle fleet. The number of vehicle miles traveled by light-duty motor vehicles (passenger cars and light-duty trucks) increased 39 percent from 1990 to 2009, as a result of a confluence of factors including population growth, economic growth, urban sprawl, and low fuel prices over much of this period.

Industrial End-Use Sector. Industrial CO_2 emissions, resulting both directly from the combustion of fossil fuels and indirectly from the generation of electricity that is consumed by industry, accounted for 26 percent of CO_2 from fossil fuel combustion in 2009. Approximately 55 percent of these emissions resulted from direct fossil fuel combustion to produce steam and/or heat for industrial processes. The remaining emissions resulted from consuming electricity for motors, electric furnaces, ovens, lighting, and other applications. In contrast to the other end-use sectors, emissions from industry have steadily declined since 1990. This decline is due to structural changes in the U.S. economy (i.e., shifts from a manufacturing-based to a service-based economy), fuel switching, and efficiency improvements.

Residential and Commercial End-Use Sectors. The residential and commercial end-use sectors accounted for 22 and 19 percent, respectively, of CO_2 emissions from fossil fuel combustion in 2009. Both sectors relied heavily on electricity for meeting energy demands, with 70 and 77 percent, respectively, of their emissions attributable to electricity consumption for lighting, heating, cooling, and operating appliances. The remaining emissions were due to the consumption of natural gas and petroleum for heating and cooking. Emissions from these end-use sectors have increased 25 percent since 1990, due to increasing electricity consumption for lighting, heating, air conditioning, and operating appliances.

Electricity Generation. The United States relies on electricity to meet a significant portion of its energy demands. Electricity generators consumed 36 percent of U.S. energy from fossil fuels and emitted 41 percent of the CO_2 from fossil fuel combustion in 2009. The type of fuel combusted by electricity generators has a significant effect on their emissions. For example, some electricity is generated with low CO_2 emitting energy technologies, particularly non-fossil options such as nuclear, hydroelectric, or geothermal energy. However, electricity generators rely on coal for over half of their total energy requirements and accounted for 95 percent of all coal consumed for energy in the United States in 2009. Consequently, changes in electricity demand have a significant impact on coal consumption and associated CO_2 emissions.

Other significant CO_2 trends included the following:

- Carbon dioxide emissions from non-energy use of fossil fuels have increased 4.7 Tg CO_2 Eq. (4.0 percent) from 1990 through 2009. Emissions from non-energy uses of fossil fuels were 123.4 Tg CO_2 Eq. in 2009, which constituted 2.2 percent of total national CO_2 emissions, approximately the same proportion as in 1990.
- Carbon dioxide emissions from iron and steel production and metallurgical coke production decreased by 24.1 Tg CO_2 Eq. (36.6 percent) from 2008 to 2009, continuing a trend of decreasing emissions from 1990 through 2009 of 57.9 percent (57.7 Tg CO_2 Eq.). This decline is due to the restructuring of the industry, technological improvements, and increased scrap utilization.
- In 2009, CO_2 emissions from cement production decreased by 11.5 Tg CO_2 Eq. (28.4 percent) from 2008. After decreasing in 1991 by two percent from 1990 levels, cement production emissions grew every year through 2006; emissions decreased in the last three years. Overall, from 1990 to 2009, emissions from cement production decreased by 12.8 percent, a decrease of 4.3 Tg CO_2 Eq.
- Net CO_2 uptake from Land Use, Land-Use Change, and Forestry increased by 153.5 Tg CO_2 Eq. (17.8 percent) from 1990 through 2009. This increase was primarily due to an increase in the rate of net carbon accumulation in forest carbon stocks, particularly in aboveground and belowground tree biomass, and harvested wood pools. Annual

carbon accumulation in landfilled yard trimmings and food scraps slowed over this period, while the rate of carbon accumulation in urban trees increased.

Methane Emissions

Methane (CH_4) is more than 20 times as effective as CO_2 at trapping heat in the atmosphere (IPCC 1996). Over the last two hundred and fifty years, the concentration of CH_4 in the atmosphere increased by 148 percent (IPCC 2007). Anthropogenic sources of CH_4 include natural gas and petroleum systems, agricultural activities, landfills, coal mining, wastewater treatment, stationary and mobile combustion, and certain industrial processes (see Figure ES-8).

Figure ES-8



Some significant trends in U.S. emissions of CH_4 include the following:

- In 2009, CH_4 emissions from coal mining were 71.0 Tg CO_2 Eq., a 3.9 Tg CO_2 Eq. (5.8 percent) increase over 2008 emission levels. The overall decline of 13.0 Tg CO_2 Eq. (15.5 percent) from 1990 results from the mining of less gassy coal from underground mines and the increased use of CH_4 collected from degasification systems.
- Natural gas systems were the largest anthropogenic source category of CH_4 emissions in the United States in 2009 with 221.2 Tg CO_2 Eq. of CH_4 emitted into the atmosphere. Those emissions have increased by 31.4 Tg CO_2 Eq. (16.6 percent) since 1990. Methane emissions from this source increased 4 percent from 2008 to 2009 due to an increase in production and production wells.
- Enteric Fermentation is the second largest anthropogenic source of CH_4 emissions in the United States. In 2009, enteric fermentation CH_4 emissions were 139.8 Tg CO_2 Eq. (20 percent of total CH_4 emissions), which represents an increase of 7.7 Tg CO_2 Eq. (5.8 percent) since 1990.
- Methane emissions from manure management increased by 55.9 percent since 1990, from 31.7 Tg CO_2 Eq. in 1990 to 49.5 Tg CO_2 Eq. in 2009. The majority of this increase was from swine and dairy cow manure, since the general trend in manure management is one of increasing use of liquid systems, which tends to produce greater CH_4 emissions. The increase in liquid systems is the combined result of a shift to larger facilities, and to facilities in the West and Southwest, all of which tend to use liquid systems. Also, new regulations limiting the application of manure nutrients have shifted manure management practices at smaller dairies from daily spread to manure managed and stored on site.
- Landfills are the third largest anthropogenic source of CH_4 emissions in the United States, accounting for 17 percent of total CH_4 emissions (117.5 Tg CO_2 Eq.) in 2009. From 1990 to 2009, CH_4 emissions from landfills decreased by 29.9 Tg CO_2 Eq. (20 percent), with small increases occurring in some interim years. This downward trend in overall

emissions is the result of increases in the amount of landfill gas collected and combusted,[14] which has more than offset the additional CH_4 emissions resulting from an increase in the amount of municipal solid waste landfilled.

Nitrous Oxide Emissions

Nitrous oxide is produced by biological processes that occur in soil and water and by a variety of anthropogenic activities in the agricultural, energy-related, industrial, and waste management fields. While total N_2O emissions are much lower than CO_2 emissions, N_2O is approximately 300 times more powerful than CO_2 at trapping heat in the atmosphere (IPCC 1996). Since 1750, the global atmospheric concentration of N_2O has risen by approximately 18 percent (IPCC 2007). The main anthropogenic activities producing N_2O in the United States are agricultural soil management, fuel combustion in motor vehicles, manure management, nitric acid production and stationary fuel combustion, (see Figure ES-9).

Figure ES-9



Some significant trends in U.S. emissions of N_2O include the following:

- In 2009, N_2O emissions from mobile combustion were 23.9 Tg CO_2 Eq. (approximately 8.1 percent of U.S. N_2O emissions). From 1990 to 2009, N_2O emissions from mobile combustion decreased by 45.6 percent. However, from 1990 to 1998 emissions increased by 25.6 percent, due to control technologies that reduced NO_x emissions while increasing N_2O emissions. Since 1998, newer control technologies have led to an overall decline in N_2O from this source.
- Nitrous oxide emissions from adipic acid production were 1.9 Tg CO_2 Eq. in 2009, and have decreased significantly since 1996 from the widespread installation of pollution control measures. Emissions from adipic acid production have decreased by 87.7 percent since 1990, and emissions from adipic acid production have remained consistently lower than pre-1996 levels since 1998.
- Agricultural soils accounted for approximately 69.2 percent of N_2O emissions in the United States in 2009. Estimated emissions from this source in 2009 were 204.6 Tg CO_2 Eq. Annual N_2O emissions from agricultural soils fluctuated between 1990 and 2009, although overall emissions were 3.4 percent higher in 2009 than in 1990.

[14] The CO_2 produced from combusted landfill CH_4 at landfills is not counted in national inventories as it is considered part of the natural C cycle of decomposition.

HFC, PFC, and SF_6 Emissions

HFCs and PFCs are families of synthetic chemicals that are used as alternatives to ODS, which are being phased out under the *Montreal Protocol* and Clean Air Act Amendments of 1990. HFCs and PFCs do not deplete the stratospheric ozone layer, and are therefore acceptable alternatives under the *Montreal Protocol*.

These compounds, however, along with SF_6, are potent greenhouse gases. In addition to having high global warming potentials, SF_6 and PFCs have extremely long atmospheric lifetimes, resulting in their essentially irreversible accumulation in the atmosphere once emitted. Sulfur hexafluoride is the most potent greenhouse gas the IPCC has evaluated (IPCC 1996).

Other emissive sources of these gases include electrical transmission and distribution systems, HCFC-22 production, semiconductor manufacturing, aluminum production, and magnesium production and processing (see Figure ES-10).

Figure ES-10

2009 Sources of HFCs, PFCs, and SF_6 Emissions



Some significant trends in U.S. HFC, PFC, and SF_6 emissions include the following:

- Emissions resulting from the substitution of ODS (e.g., CFCs) have been consistently increasing, from small amounts in 1990 to 120.0 Tg CO_2 Eq. in 2009. Emissions from ODS substitutes are both the largest and the fastest growing source of HFC, PFC, and SF_6 emissions. These emissions have been increasing as phase-outs required under the *Montreal Protocol* come into effect, especially after 1994, when full market penetration was made for the first generation of new technologies featuring ODS substitutes.
- HFC emissions from the production of HCFC-22 decreased by 85.2 percent (31.0 Tg CO_2 Eq.) from 1990 through 2009, due to a steady decline in the emission rate of HFC-23 (i.e., the amount of HFC-23 emitted per kilogram of HCFC-22 manufactured) and the use of thermal oxidation at some plants to reduce HFC-23 emissions.
- Sulfur hexafluoride emissions from electric power transmission and distribution systems decreased by 54.8 percent (15.6 Tg CO_2 Eq.) from 1990 to 2009, primarily because of higher purchase prices for SF_6 and efforts by industry to reduce emissions.
- PFC emissions from aluminum production decreased by 91.5 percent (17.0 Tg CO_2 Eq.) from 1990 to 2009, due to both industry emission reduction efforts and lower domestic aluminum production.

ES.3. Overview of Sector Emissions and Trends

In accordance with the *Revised 1996 IPCC Guidelines for National Greenhouse Gas Inventories* (IPCC/UNEP/OECD/IEA 1997), and the 2003 UNFCCC Guidelines on Reporting and Review (UNFCCC 2003), Figure ES-11 and Table ES-4 aggregate emissions and sinks by these chapters. Emissions of all gases can be summed from each source category from IPCC guidance. Over the twenty-year period of 1990 to 2009, total emissions in the Energy and Agriculture sectors grew by 463.3 Tg CO_2 Eq. (9 percent), and 35.7 Tg CO_2 Eq. (9 percent), respectively. Emissions decreased in the Industrial Processes, Waste, and Solvent and Other Product Use sectors by 32.9 Tg CO_2 Eq. (10 percent), 24.7 Tg CO_2 Eq. (14 percent) and less than 0.1 Tg CO_2 Eq. (0.4 percent), respectively. Over the same period, estimates of net C sequestration in the Land Use, Land-Use Change, and Forestry sector (magnitude of emissions plus CO_2 flux from all LULUCF source categories) increased by 143.5 Tg CO_2 Eq. (17 percent).

Figure ES-11

U.S. Greenhouse Gas Emissions and Sinks by Chapter/IPCC Sector



Note: Relatively smaller amounts of GWP-weighted emissions are also emitted from the Solvent and Other Product Use Sectors.

Table ES-4: Recent Trends in U.S. Greenhouse Gas Emissions and Sinks by Chapter/IPCC Sector (Tg or million metric tons CO_2 Eq.)

Chapter/IPCC Sector	1990	2000	2005	2006	2007	2008	2009
Energy	5,287.8	6,168.0	6,282.8	6,210.2	6,290.7	6,116.6	5,751.1
Industrial Processes	315.8	348.8	334.1	339.4	350.9	331.7	282.9
Solvent and Other Product Use	4.4	4.9	4.4	4.4	4.4	4.4	4.4
Agriculture	383.6	410.6	418.8	418.8	425.8	426.3	419.3
Land Use, Land-Use Change, and Forestry (Emissions)	15.0	36.3	28.6	49.8	47.5	33.2	25.0
Waste	175.2	143.9	144.9	144.4	144.1	149.0	150.5
Total Emissions	**6,181.8**	**7,112.7**	**7,213.5**	**7,166.9**	**7,263.4**	**7,061.1**	**6,633.2**
Net CO_2 Flux from Land Use, Land-Use Change, and Forestry (Sinks)[a]	(861.5)	(576.6)	(1,056.5)	(1,064.3)	(1,060.9)	(1,040.5)	(1,015.1)
Net Emissions (Sources and Sinks)	**5,320.3**	**6,536.1**	**6,157.1**	**6,102.6**	**6,202.5**	**6,020.7**	**5,618.2**

[a] The net CO_2 flux total includes both emissions and sequestration, and constitutes a sink in the United States. Sinks are only included in net emissions total.

Note: Totals may not sum due to independent rounding. Parentheses indicate negative values or sequestration.

Energy

The Energy chapter contains emissions of all greenhouse gases resulting from stationary and mobile energy activities including fuel combustion and fugitive fuel emissions. Energy-related activities, primarily fossil fuel combustion, accounted for the vast majority of U.S. CO_2 emissions for the period of 1990 through 2009. In 2009, approximately 83 percent of the energy consumed in the United States (on a Btu basis) was produced through the combustion of fossil fuels. The remaining 17 percent came from other energy sources such as hydropower, biomass, nuclear, wind, and solar energy (see Figure ES-12). Energy-related activities are also responsible for CH_4 and N_2O emissions (49 percent and 13 percent of total U.S. emissions of each gas, respectively). Overall, emission sources in the Energy chapter account for a combined 87 percent of total U.S. greenhouse gas emissions in 2009.

Figure ES-12

2009 U.S. Energy Consumption by Energy Source



Industrial Processes

The Industrial Processes chapter contains byproduct or fugitive emissions of greenhouse gases from industrial processes not directly related to energy activities such as fossil fuel combustion. For example, industrial processes can chemically transform raw materials, which often release waste gases such as CO_2, CH_4, and N_2O. These processes include iron and steel production and metallurgical coke production, cement production, ammonia production and urea consumption, lime production, limestone and dolomite use (e.g., flux stone, flue gas desulfurization, and glass manufacturing), soda ash production and consumption, titanium dioxide production, phosphoric acid production, ferroalloy production, CO_2 consumption, silicon carbide production and consumption, aluminum production, petrochemical production, nitric acid production, adipic acid production, lead production, and zinc production. Additionally, emissions from industrial processes release HFCs, PFCs, and SF_6. Overall, emission sources in the Industrial Process chapter account for 4 percent of U.S. greenhouse gas emissions in 2009.

Solvent and Other Product Use

The Solvent and Other Product Use chapter contains greenhouse gas emissions that are produced as a byproduct of various solvent and other product uses. In the United States, emissions from N_2O from product uses, the only source of greenhouse gas emissions from this sector, accounted for about 0.1 percent of total U.S. anthropogenic greenhouse gas emissions on a carbon equivalent basis in 2009.

Agriculture

The Agriculture chapter contains anthropogenic emissions from agricultural activities (except fuel combustion, which is addressed in the Energy chapter, and agricultural CO_2 fluxes, which are addressed in the Land Use, Land-Use Change, and Forestry Chapter). Agricultural activities contribute directly to emissions of greenhouse gases through a variety of processes, including the following source categories: enteric fermentation in domestic livestock, livestock manure management, rice cultivation, agricultural soil management, and field burning of agricultural residues. CH_4 and N_2O were the primary greenhouse gases emitted by agricultural activities. Methane emissions from enteric fermentation and manure management represented 20 percent and 7 percent of total CH_4 emissions from anthropogenic activities, respectively, in 2009.

Agricultural soil management activities such as fertilizer application and other cropping practices were the largest source of U.S. N_2O emissions in 2009, accounting for 69 percent. In 2009, emission sources accounted for in the Agriculture chapter were responsible for 6.3 percent of total U.S. greenhouse gas emissions.

Land Use, Land-Use Change, and Forestry

The Land Use, Land-Use Change, and Forestry chapter contains emissions of CH_4 and N_2O, and emissions and removals of CO_2 from forest management, other land-use activities, and land-use change. Forest management practices, tree planting in urban areas, the management of agricultural soils, and the landfilling of yard trimmings and food scraps resulted in a net uptake (sequestration) of C in the United States. Forests (including vegetation, soils, and harvested wood) accounted for 85 percent of total 2009 net CO_2 flux, urban trees accounted for 9 percent, mineral and organic soil carbon stock changes accounted for 4 percent, and landfilled yard trimmings and food scraps accounted for 1 percent of the total net flux in 2009. The net forest sequestration is a result of net forest growth and increasing forest area, as well as a net accumulation of carbon stocks in harvested wood pools. The net sequestration in urban forests is a result of net tree growth in these areas. In agricultural soils, mineral and organic soils sequester approximately 5.5 times as much C as is emitted from these soils through liming and urea fertilization. The mineral soil C sequestration is largely due to the conversion of cropland to permanent pastures and hay production, a reduction in summer fallow areas in semi-arid areas, an increase in the adoption of conservation tillage practices, and an increase in the amounts of organic fertilizers (i.e., manure and sewage sludge) applied to agriculture lands. The landfilled yard trimmings and food scraps net sequestration is due to the long-term accumulation of yard trimming carbon and food scraps in landfills.

Land use, land-use change, and forestry activities in 2009 resulted in a net C sequestration of 1,015.1 Tg CO_2 Eq. (Table ES-5). This represents an offset of 18 percent of total U.S. CO_2 emissions, or 15 percent of total greenhouse gas emissions in 2009. Between 1990 and 2009, total land use, land-use change, and forestry net C flux resulted in a 17.8 percent increase in CO_2 sequestration, primarily due to an increase in the rate of net C accumulation in forest C stocks, particularly in aboveground and belowground tree biomass, and harvested wood pools. Annual C accumulation in landfilled yard trimmings and food scraps slowed over this period, while the rate of annual C accumulation increased in urban trees.

Table ES-5: Net CO_2 Flux from Land Use, Land-Use Change, and Forestry (Tg or million metric tons CO_2 Eq.)

Sink Category	1990	2000	2005	2006	2007	2008	2009
Forest Land Remaining Forest Land	(681.1)	(378.3)	(911.5)	(917.5)	(911.9)	(891.0)	(863.1)
Cropland Remaining Cropland	(29.4)	(30.2)	(18.3)	(19.1)	(19.7)	(18.1)	(17.4)
Land Converted to Cropland	2.2	2.4	5.9	5.9	5.9	5.9	5.9
Grassland Remaining Grassland	(52.2)	(52.6)	(8.9)	(8.8)	(8.6)	(8.5)	(8.3)
Land Converted to Grassland	(19.8)	(27.2)	(24.4)	(24.2)	(24.0)	(23.8)	(23.6)
Settlements Remaining Settlements	(57.1)	(77.5)	(87.8)	(89.8)	(91.9)	(93.9)	(95.9)
Other (Landfilled Yard Trimmings and Food Scraps)	(24.2)	(13.2)	(11.5)	(11.0)	(10.9)	(11.2)	(12.6)
Total	**(861.5)**	**(576.6)**	**(1,056.5)**	**(1,064.3)**	**(1,060.9)**	**(1,040.5)**	**(1,015.1)**

Note: Totals may not sum due to independent rounding. Parentheses indicate net sequestration.

Emissions from Land Use, Land-Use Change, and Forestry are shown in Table ES-6. The application of crushed limestone and dolomite to managed land (i.e., liming of agricultural soils) and urea fertilization resulted in CO_2 emissions of 7.8 Tg CO_2 Eq. in 2009, an increase of 11 percent relative to 1990. The application of synthetic fertilizers to forest and settlement soils in 2009 resulted in direct N_2O emissions of 1.9 Tg CO_2 Eq. Direct N_2O emissions from fertilizer application to forest soils have increased by 455 percent since 1990, but still account for a relatively small portion of overall emissions. Additionally, direct N_2O emissions from fertilizer application to settlement soils increased by 55 percent since 1990. Forest

fires resulted in CH_4 emissions of 7.8 Tg CO_2 Eq., and in N_2O emissions of 6.4 Tg CO_2 Eq. in 2009. Carbon dioxide and N_2O emissions from peatlands totaled 1.1 Tg CO_2 Eq. and less than 0.01 Tg CO_2 Eq. in 2009, respectively.

Table ES-6: Emissions from Land Use, Land-Use Change, and Forestry (Tg or million metric tons CO_2 Eq.)

Source Category	1990	2000	2005	2006	2007	2008	2009
CO_2	**8.1**	**8.8**	**8.9**	**8.8**	**9.2**	**9.6**	**8.9**
Cropland Remaining Cropland: Liming of Agricultural Soils	4.7	4.3	4.3	4.2	4.5	5.0	4.2
Cropland Remaining Cropland: Urea Fertilization	2.4	3.2	3.5	3.7	3.7	3.6	3.6
Wetlands Remaining Wetlands: Peatlands Remaining Peatlands	1.0	1.2	1.1	0.9	1.0	1.0	1.1
CH_4	**3.2**	**14.3**	**9.8**	**21.6**	**20.0**	**11.9**	**7.8**
Forest Land Remaining Forest Land: Forest Fires	3.2	14.3	9.8	21.6	20.0	11.9	7.8
N_2O	**3.7**	**13.2**	**9.8**	**19.5**	**18.3**	**11.6**	**8.3**
Forest Land Remaining Forest Land: Forest Fires	2.6	11.7	8.0	17.6	16.3	9.8	6.4
Forest Land Remaining Forest Land: Forest Soils	0.1	0.4	0.4	0.4	0.4	0.4	0.4
Settlements Remaining Settlements: Settlement Soils	1.0	1.1	1.5	1.5	1.6	1.5	1.5
Wetlands Remaining Wetlands: Peatlands Remaining Peatlands	+	+	+	+	+	+	+
Total	**15.0**	**36.3**	**28.6**	**49.8**	**47.5**	**33.2**	**25.0**

+ Less than 0.05 Tg CO_2 Eq.
Note: Totals may not sum due to independent rounding.

Waste

The Waste chapter contains emissions from waste management activities (except incineration of waste, which is addressed in the Energy chapter). Landfills were the largest source of anthropogenic greenhouse gas emissions in the Waste chapter, accounting for just over 78 percent of this chapter's emissions, and 17 percent of total U.S. CH_4 emissions.[15] Additionally, wastewater treatment accounts for 20 percent of Waste emissions, 4 percent of U.S. CH_4 emissions, and 2 percent of U.S. N_2O emissions. Emissions of CH_4 and N_2O from composting are also accounted for in this chapter; generating emissions of 1.7 Tg CO_2 Eq. and 1.8 Tg CO_2 Eq., respectively. Overall, emission sources accounted for in the Waste chapter generated 2.3 percent of total U.S. greenhouse gas emissions in 2009.

ES.4. Other Information

Emissions by Economic Sector

Throughout the Inventory of U.S. Greenhouse Gas Emissions and Sinks report, emission estimates are grouped into six sectors (i.e., chapters) defined by the IPCC: Energy; Industrial Processes; Solvent Use; Agriculture; Land Use, Land-Use Change, and Forestry; and Waste. While it is important to use this characterization for consistency with UNFCCC reporting guidelines, it is also useful to allocate emissions into more commonly used sectoral categories. This section reports emissions by the following economic sectors: Residential, Commercial, Industry, Transportation, Electricity Generation, Agriculture, and U.S. Territories.

Table ES-7 summarizes emissions from each of these sectors, and Figure ES-13 shows the trend in emissions by sector from 1990 to 2009.

[15] Landfills also store carbon, due to incomplete degradation of organic materials such as wood products and yard trimmings, as described in the Land-Use, Land-Use Change, and Forestry chapter of the inventory report.

Table ES-7: U.S. Greenhouse Gas Emissions Allocated to Economic Sectors (Tg or million metric tons CO_2 Eq.)

Implied Sectors	1990	2000	2005	2006	2007	2008	2009
Electric Power Industry	1,868.9	2,337.6	2,444.6	2,388.2	2,454.0	2,400.7	2,193.0
Transportation	1,545.2	1,932.3	2,017.4	1,994.4	2,003.8	1,890.7	1,812.4
Industry	1,564.4	1,544.0	1,441.9	1,497.3	1,483.0	1,446.9	1,322.7
Agriculture	429.0	485.1	493.2	516.7	520.7	503.9	490.0
Commercial	395.5	381.4	387.2	375.2	389.6	403.5	409.5
Residential	345.1	386.2	371.0	335.8	358.9	367.1	360.1
U.S. Territories	33.7	46.0	58.2	59.3	53.5	48.4	45.5
Total Emissions	**6,181.8**	**7,112.7**	**7,213.5**	**7,166.9**	**7,263.4**	**7,061.1**	**6,633.2**
Land Use, Land-Use Change, and Forestry (Sinks)	(861.5)	(576.6)	(1,056.5)	(1,064.3)	(1,060.9)	(1,040.5)	(1,015.1)
Net Emissions (Sources and Sinks)	**5,320.3**	**6,536.1**	**6,157.1**	**6,102.6**	**6,202.5**	**6,020.7**	**5,618.2**

Note: Totals may not sum due to independent rounding. Emissions include CO_2, CH_4, N_2O, HFCs, PFCs, and SF_6.
See Table 2-12 of the inventory report for more detailed data.

Figure ES-13

Emissions Allocated to Economic Sectors



Note: Does not include U.S. Territories.

Using this categorization, emissions from electricity generation accounted for the largest portion (33 percent) of U.S. greenhouse gas emissions in 2009. Transportation activities, in aggregate, accounted for the second largest portion (27 percent), while emissions from industry accounted for the third largest portion (20 percent) of U.S. greenhouse gas emissions in 2009. In contrast to electricity generation and transportation, emissions from industry have in general declined over the past decade. The long-term decline in these emissions has been due to structural changes in the U.S. economy (i.e., shifts from a manufacturing-based to a service-based economy), fuel switching, and energy efficiency improvements. The remaining 20 percent of U.S. greenhouse gas emissions were contributed by, in order of importance, the agriculture, commercial, and residential sectors, plus emissions from U.S. territories. Activities related to agriculture accounted for 7 percent of U.S. emissions; unlike other economic sectors, agricultural sector emissions were dominated by N_2O emissions from agricultural soil management and CH_4 emissions from enteric fermentation. The commercial sector accounted for 6 percent of emissions while the residential sector accounted for 5 percent of emissions and U.S. territories accounted for 1 percent of emissions; emissions from these sectors primarily consisted of CO_2 emissions from fossil fuel combustion.

Carbon dioxide was also emitted and sequestered by a variety of activities related to forest management practices, tree planting in urban areas, the management of agricultural soils, and landfilling of yard trimmings.

Electricity is ultimately consumed in the economic sectors described above. Table ES-8 presents greenhouse gas emissions from economic sectors with emissions related to electricity generation distributed into end-use categories (i.e., emissions from electricity generation are allocated to the economic sectors in which the electricity is consumed). To distribute electricity emissions among end-use sectors, emissions from the source categories assigned to electricity generation were allocated to the residential, commercial, industry, transportation, and agriculture economic sectors according to retail

sales of electricity.[16] These source categories include CO_2 from fossil fuel combustion and the use of limestone and dolomite for flue gas desulfurization, CO_2 and N_2O from incineration of waste, CH_4 and N_2O from stationary sources, and SF_6 from electrical transmission and distribution systems.

Table ES-8: U.S. Greenhouse Gas Emissions by Economic Sector with Electricity-Related Emissions Distributed (Tg or million metric tons CO_2 Eq.)

Implied Sectors	1990	2000	2005	2006	2007	2008	2009
Industry	2,238.3	2,314.4	2,162.5	2,194.6	2,192.9	2,146.5	1,910.9
Transportation	1,548.3	1,935.8	2,022.2	1,999.0	2,008.9	1,895.5	1,816.9
Commercial	947.7	1,135.8	1,205.1	1,188.5	1,225.3	1,224.5	1,184.9
Residential	953.8	1,162.2	1,242.9	1,181.5	1,229.6	1,215.1	1,158.9
Agriculture	460.0	518.4	522.7	544.1	553.2	531.1	516.0
U.S. Territories	33.7	46.0	58.2	59.3	53.5	48.4	45.5
Total Emissions	**6,181.8**	**7,112.7**	**7,213.5**	**7,166.9**	**7,263.4**	**7,061.1**	**6,633.2**
Land Use, Land-Use Change, and Forestry (Sinks)	(861.5)	(576.6)	(1,056.5)	(1,064.3)	(1,060.9)	(1,040.5)	(1,015.1)
Net Emissions (Sources and Sinks)	**5,320.3**	**6,536.1**	**6,157.1**	**6,102.6**	**6,202.5**	**6,020.7**	**5,618.2**

See Table 2-14 of the inventory report for more detailed data.

When emissions from electricity are distributed among these sectors, industrial activities account for the largest share of U.S. greenhouse gas emissions (29 percent) in 2009. Transportation is the second largest contributor to total U.S. emissions (28 percent). The commercial and residential sectors contributed the next largest shares of total U.S. greenhouse gas emissions in 2009. Emissions from these sectors increase substantially when emissions from electricity are included, due to their relatively large share of electricity consumption (e.g., lighting, appliances, etc.). In all sectors except agriculture, CO_2 accounts for more than 80 percent of greenhouse gas emissions, primarily from the combustion of fossil fuels. Figure ES-14 shows the trend in these emissions by sector from 1990 to 2009.

Figure ES-14

U.S. Emissions with Electricity Distributed to Economic Sectors



Note: Does not include U.S. Territories.

[16] Emissions were not distributed to U.S. territories, since the electricity generation sector only includes emissions related to the generation of electricity in the 50 states and the District of Columbia.

Box ES-2: Recent Trends in Various U.S. Greenhouse Gas Emissions-Related Data

Total emissions can be compared to other economic and social indices to highlight changes over time. These comparisons include: (1) emissions per unit of aggregate energy consumption, because energy-related activities are the largest sources of emissions; (2) emissions per unit of fossil fuel consumption, because almost all energy-related emissions involve the combustion of fossil fuels; (3) emissions per unit of electricity consumption, because the electric power industry—utilities and nonutilities combined—was the largest source of U.S. greenhouse gas emissions in 2009; (4) emissions per unit of total gross domestic product as a measure of national economic activity; and (5) emissions per capita.

Table ES-9 provides data on various statistics related to U.S. greenhouse gas emissions normalized to 1990 as a baseline year. Greenhouse gas emissions in the United States have grown at an average annual rate of 0.4 percent since 1990. This rate is slightly slower than that for total energy and for fossil fuel consumption, and much slower than that for electricity consumption, overall gross domestic product and national population (see Figure ES-15).

Table ES-9: Recent Trends in Various U.S. Data (Index 1990 = 100)

Variable	1990	2000	2005	2006	2007	2008	2009	Growth Rate[a]
GDP [b]	100	140	157	162	165	165	160	2.5%
Electricity Consumption [c]	100	127	134	135	138	138	132	1.5%
Fossil Fuel Consumption [c]	100	117	119	117	119	116	108	0.5%
Energy Consumption [c]	100	116	118	118	120	118	112	0.6%
Population [d]	100	113	118	120	121	122	123	1.1%
Greenhouse Gas Emissions [e]	100	115	117	116	117	114	107	0.4%

[a] Average annual growth rate
[b] Gross Domestic Product in chained 2005 dollars (BEA 2010)
[c] Energy content-weighted values (EIA 2010b)
[d] U.S. Census Bureau (2010)
[e] GWP-weighted values

Figure ES-15

U.S. Greenhouse Gas Emissions Per Capita and Per Dollar of Gross Domestic Product



Source: BEA (2010), U.S. Census Bureau (2010), and emission estimates in the inventory report.

Indirect Greenhouse Gases (CO, NO_x, NMVOCs, and SO_2)

The reporting requirements of the UNFCCC[17] request that information be provided on indirect greenhouse gases, which include CO, NO_x, NMVOCs, and SO_2. These gases do not have a direct global warming effect, but indirectly affect terrestrial radiation absorption by influencing the formation and destruction of tropospheric and stratospheric ozone, or, in the case of SO_2, by affecting the absorptive characteristics of the atmosphere. Additionally, some of these gases may react with other chemical compounds in the atmosphere to form compounds that are greenhouse gases.

Since 1970, the United States has published estimates of annual emissions of CO, NO_x, NMVOCs, and SO_2 (EPA 2010, EPA 2009),[18] which are regulated under the Clean Air Act. Table ES-10 shows that fuel combustion accounts for the majority of emissions of these indirect greenhouse gases. Industrial processes—such as the manufacture of chemical and allied products, metals processing, and industrial uses of solvents—are also significant sources of CO, NO_x, and NMVOCs.

Table ES-10: Emissions of NO_x, CO, NMVOCs, and SO_2 (Gg)

Gas/Activity	1990	2000	2005	2006	2007	2008	2009
NO_x	**21,707**	**19,116**	**15,900**	**15,039**	**14,380**	**13,547**	**11,468**
Mobile Fossil Fuel Combustion	10,862	10,199	9,012	8,488	7,965	7,441	6,206
Stationary Fossil Fuel Combustion	10,023	8,053	5,858	5,545	5,432	5,148	4,159
Industrial Processes	591	626	569	553	537	520	568
Oil and Gas Activities	139	111	321	319	318	318	393
Incineration of Waste	82	114	129	121	114	106	128
Agricultural Burning	8	8	6	7	8	8	8
Solvent Use	1	3	3	4	4	4	3
Waste	0	2	2	2	2	2	2
CO	**130,038**	**92,243**	**70,809**	**67,238**	**63,625**	**60,039**	**51,452**
Mobile Fossil Fuel Combustion	119,360	83,559	62,692	58,972	55,253	51,533	43,355
Stationary Fossil Fuel Combustion	5,000	4,340	4,649	4,695	4,744	4,792	4,543
Industrial Processes	4,125	2,216	1,555	1,597	1,640	1,682	1,549
Incineration of Waste	978	1,670	1,403	1,412	1,421	1,430	1,403
Agricultural Burning	268	259	184	233	237	270	247
Oil and Gas Activities	302	146	318	319	320	322	345
Waste	1	8	7	7	7	7	7
Solvent Use	5	45	2	2	2	2	2
NMVOCs	**20,930**	**15,227**	**13,761**	**13,594**	**13,423**	**13,254**	**9,313**
Mobile Fossil Fuel Combustion	10,932	7,229	6,330	6,037	5,742	5,447	4,151
Solvent Use	5,216	4,384	3,851	3,846	3,839	3,834	2,583
Industrial Processes	2,422	1,773	1,997	1,933	1,869	1,804	1,322
Stationary Fossil Fuel Combustion	912	1,077	716	918	1,120	1,321	424
Oil and Gas Activities	554	388	510	510	509	509	599
Incineration of Waste	222	257	241	238	234	230	159
Waste	673	119	114	113	111	109	76
Agricultural Burning	NA	NA	NA	NA	NA	NA	NA
SO_2	**20,935**	**14,830**	**13,466**	**12,388**	**11,799**	**10,368**	**8,599**
Stationary Fossil Fuel Combustion	18,407	12,849	11,541	10,612	10,172	8,891	7,167
Industrial Processes	1,307	1,031	831	818	807	795	798
Mobile Fossil Fuel Combustion	793	632	889	750	611	472	455
Oil and Gas Activities	390	287	181	182	184	187	154
Incineration of Waste	38	29	24	24	24	23	24
Waste	0	1	1	1	1	1	1
Solvent Use	0	1	0	0	0	0	0
Agricultural Burning	NA	NA	NA	NA	NA	NA	NA

NA (Not Available)
Note: Totals may not sum due to independent rounding.
Source: (EPA 2010, EPA 2009) except for estimates from field burning of agricultural residues.

[17] See <http://unfccc.int/resource/docs/cop8/08.pdf>.

[18] NO_x and CO emission estimates from field burning of agricultural residues were estimated separately, and therefore not taken from EPA (2008).

Key Categories

The *2006 IPCC Guidelines for National Greenhouse Gas Inventories* (IPCC 2006) defines a key category as a "[source or sink category] that is prioritized within the national inventory system because its estimate has a significant influence on a country's total inventory of direct greenhouse gases in terms of the absolute level of emissions, the trend in emissions, or both."[19] By definition, key categories are sources or sinks that have the greatest contribution to the absolute overall level of national emissions in any of the years covered by the time series. In addition, when an entire time series of emission estimates is prepared, a thorough investigation of key categories must also account for the influence of trends of individual source and sink categories. Finally, a qualitative evaluation of key categories should be performed, in order to capture any key categories that were not identified in either of the quantitative analyses.

Figure ES-16

2009 Key Categories



Note: For a complete discussion of the key category analysis, see Annex 1 of the inventory report. Darker bars indicate a Tier 1 level assessment key category. Lighter bars indicate a Tier 2 level assessment key category.

Figure ES-16 presents 2009 emission estimates for the key categories as defined by a level analysis (i.e., the contribution of each source or sink category to the total inventory level). The UNFCCC reporting guidelines request that key category analyses be reported at an appropriate level of disaggregation, which may lead to source and sink category names which differ from those used elsewhere in the inventory report. For more information regarding key categories, see section 1.5 and Annex 1.

[19] See Chapter 7 "Methodological Choice and Recalculation" in IPCC (2000). <http://www.ipcc-nggip.iges.or.jp/public/gp/gpgaum.htm>.

Quality Assurance and Quality Control (QA/QC)

The United States seeks to continually improve the quality, transparency, and credibility of the Inventory of U.S. Greenhouse Gas Emissions and Sinks. To assist in these efforts, the United States implemented a systematic approach to QA/QC. While QA/QC has always been an integral part of the U.S. national system for inventory development, the procedures followed for the current inventory have been formalized in accordance with the QA/QC plan and the UNFCCC reporting guidelines.

Uncertainty Analysis of Emission Estimates

While the current U.S. emissions inventory provides a solid foundation for the development of a more detailed and comprehensive national inventory, there are uncertainties associated with the emission estimates. Some of the current estimates, such as those for CO_2 emissions from energy-related activities and cement processing, are considered to have low uncertainties. For some other categories of emissions, however, a lack of data or an incomplete understanding of how emissions are generated increases the uncertainty associated with the estimates presented. Acquiring a better understanding of the uncertainty associated with inventory estimates is an important step in helping to prioritize future work and improve the overall quality of the inventory report. Recognizing the benefit of conducting an uncertainty analysis, the UNFCCC reporting guidelines follow the recommendations of the IPCC Good Practice Guidance (IPCC 2000) and require that countries provide single estimates of uncertainty for source and sink categories.

Currently, a qualitative discussion of uncertainty is presented for all source and sink categories. Within the discussion of each emission source, specific factors affecting the uncertainty surrounding the estimates are discussed. Most sources also contain a quantitative uncertainty assessment, in accordance with UNFCCC reporting guidelines.

Box ES-3: Recalculations of Inventory Estimates

Each year, emission and sink estimates are recalculated and revised for all years in the Inventory of U.S. Greenhouse Gas Emissions and Sinks, as attempts are made to improve both the analyses themselves, through the use of better methods or data, and the overall usefulness of the inventory report. In this effort, the United States follows the 2006 IPCC Guidelines (IPCC 2006), which states, “Both methodological changes and refinements over time are an essential part of improving inventory quality. It is good practice to change or refine methods” when: available data have changed; the previously used method is not consistent with the IPCC guidelines for that category; a category has become key; the previously used method is insufficient to reflect mitigation activities in a transparent manner; the capacity for inventory preparation has increased; new inventory methods become available; and for correction of errors.” In general, recalculations are made to the U.S. greenhouse gas emission estimates either to incorporate new methodologies or, most commonly, to update recent historical data.

In each inventory report, the results of all methodology changes and historical data updates are presented in the "Recalculations and Improvements" chapter; detailed descriptions of each recalculation are contained within each source's description contained in the report, if applicable. In general, when methodological changes have been implemented, the entire time series (in the case of the most recent inventory report, 1990 through 2009) has been recalculated to reflect the change, per the 2006 IPCC Guidelines (IPCC 2006). Changes in historical data are generally the result of changes in statistical data supplied by other agencies. References for the data are provided for additional information.

References

BEA (2010) *2009 Comprehensive Revision of the National Income and Product Accounts: Current-dollar and "real" GDP, 1929–2009*. Bureau of Economic Analysis (BEA), U.S. Department of Commerce, Washington, DC. July 29, 2010. Available online at < http://www.bea.gov/national/index.htm#gdp >.

EIA (2010) Supplemental Tables on Petroleum Product detail. *Monthly Energy Review, September 2010,* Energy Information Administration, U.S. Department of Energy, Washington, DC. DOE/EIA-0035(2009/09).

EIA (2009) International Energy Annual 2007. Energy Information Administration (EIA), U.S. Department of Energy. Washington, DC. Updated October 2008. Available online at <http://www.eia.doe.gov/emeu/iea/carbon.html >.

EPA (2010) "2009 Average annual emissions, all criteria pollutants in MS Excel." *National Emissions Inventory (NEI) Air Pollutant Emissions Trends Data*. Office of Air Quality Planning and Standards.

EPA (2009) "1970 - 2008 Average annual emissions, all criteria pollutants in MS Excel." *National Emissions Inventory (NEI) Air Pollutant Emissions Trends Data*. Office of Air Quality Planning and Standards. Available online at <http://www.epa.gov/ttn/chief/trends/index.html>

IPCC (2007) Climate Change 2007: The Physical Science Basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change. S. Solomon , D. Qin, M. Manning, Z. Chen, M. Marquis, K.B. Averyt, M. Tignor and H.L. Miller (eds.). Cambridge University Press. Cambridge, United Kingdom 996 pp.

IPCC (2006) *2006 IPCC Guidelines for National Greenhouse Gas Inventories*. The National Greenhouse Gas Inventories Programme, The Intergovernmental Panel on Climate Change, H.S. Eggleston, L. Buendia, K. Miwa, T. Ngara, and K. Tanabe (eds.). Hayama, Kanagawa, Japan.

IPCC (2003) *Good Practice Guidance for Land Use*, Land-Use Change, and Forestry. National Greenhouse Gas Inventories Programme, The Intergovernmental Panel on Climate Change, J. Penman, et al. (eds.). Available online at <http://www.ipcc-nggip.iges.or.jp/public/gpglulucf/gpglulucf.htm>. August 13, 2004.

IPCC (2001) Climate Change 2001: The Scientific Basis. Intergovernmental Panel on Climate Change, J.T. Houghton, Y. Ding, D.J. Griggs, M. Noguer, P.J. van der Linden, X. Dai, C.A. Johnson, and K. Maskell (eds.). Cambridge University Press. Cambridge, United Kingdom.

IPCC (2000) *Good Practice Guidance and Uncertainty Management in National Greenhouse Gas Inventories*. , National Greenhouse Gas Inventories Programme, Intergovernmental Panel on Climate Change. Montreal. May 2000. IPCC-XVI/Doc. 10 (1.IV.2000).

IPCC (1996) Climate Change 1995: The Science of Climate Change. Intergovernmental Panel on Climate Change, J.T. Houghton, L.G. Meira Filho, B.A. Callander, N. Harris, A. Kattenberg, and K. Maskell. (eds.). Cambridge University Press. Cambridge, United Kingdom.

IPCC/UNEP/OECD/IEA (1997) *Revised 1996 IPCC Guidelines for National Greenhouse Gas Inventories*. Intergovernmental Panel on Climate Change, United Nations Environment Programme, Organization for Economic Co-Operation and Development, International Energy Agency. Paris, France.

UNFCCC (2003) National Communications: Greenhouse Gas Inventories from Parties included in Annex I to the Convention, UNFCCC Guidelines on Reporting and Review. Conference of the Parties, Eighth Session, New Delhi. (FCCC/CP/2002/8). March 28, 2003.

U.S. Census Bureau (2010) U.S. Census Bureau International Database (IDB). Available online at <http://www.census.gov/ipc/www/idbnew.html>. August 15, 2010.

United States
Environmental Protection Agency

EPA 430-S-11-001
April 2011
Office of Atmospheric Programs (6207J)
Washington, DC 20460

Official Business
Penalty for Private Use